
2016

Ontario Winter Wheat Performance Trials

Data collected 2012-2016

www.gocereals.ca



Conducted by the Ontario Cereal Crop Committee

Current as of September 9, 2016 (version 1)

Ontario Winter Wheat Performance Trials

This report has been prepared by the Ontario Cereal Crop Committee and contains the most recent varietal information on winter wheat that was planted in 2015 and harvested in 2016

ADDITIONAL INFORMATION

Additional information is available at www.gocereals.ca, including

- Map with area and individual locations
- Head to Head comparisons which give multi-year data
- Historical Performance reports

For more information contact:

Ellen Sparry
Trial Coordinator
C&M Seeds
519-343-2126
esparry@redwheat.com

Peter Johnson
Agronomist, Real Agriculture
519-318-2040
peter.johnson@bell.net









2016

Ontario Winter Wheat Performance Trials

Conducted by the Ontario Cereal Crop Committee ♦ www.gocereals.ca

Table #	Table	Page #
1	Cumulative Yield Index Summary - Area I & II combined	4
1a	Cumulative Yield Index Summary - Area I & II combined, INTENSIVE	5
2	Cumulative Yield Index Summary - Area I	6
2a	Cumulative Yield Index Summary - Area I INTENSIVE	7
3	Cumulative Yield Index Summary - Area II	8
3a	Cumulative Yield Index Summary - Area II INTENSIVE	9
4	Cumulative Yield Index Summary - Area III	10
5a	Varietal Characteristics - Area I	11
5b	Varietal Characteristics - Area II	12
5c	Varietal Characteristics - Area III	13
6	Cumulative Yield Index Summary - Area V	14
6a	Cumulative Yield Index Summary - Area V INTENSIVE	15
6b	Varietal Characteristics - Area V	16
7a	Ontario Winter Wheat Distributors	17
7b	Distributor Contact Information	18

**Table 1 - Ontario Winter Wheat Performance Trial; Cumulative Yield Index¹
Summary for Area I & II Combined², OCCC, August 2016**


Class ³	Variety	5 year	4 year	3 year	2 year	2016	
sww	Venture	89 ⁴	86	83	79	68	
	25W31 (awned) 				98	94	
srw	Emmit	100	99	95	93	90	
	Branson	106	106	105	107	108	
	Wave	102	101	98	98	99	
	CM614	104	104	104	105	106	
	WB425 (awned) 	104	104	103	104	105	
	CM249 (awned)	106	105	103	104	107	
	OAC Flight (awned) 	96	95	92	88	81	
	25R34 (awned)	109	109	108	108	111	
	25R40 (awned)	109	109	109	110	113	
	HY 412-SRW 	101	102	101	101	99	
	Emperor 	102	102	99	99	98	
	Marker		104	105	105	107	
	UGRC Ring (awned)			105	106	106	111
	UGRC C2-5				100	98	95
	25R46 (awned)				92	89	84
	Cruze (awned)					105	108
	DS572SRW 					102	101
	25R39					109	109
	Drew (awned)						94
	Curtis 						88
hrw	AC Morley	96	95	93	92	88	
	AC Sampson	94	94	94	93	89	
	Princeton	98	98	98	98	102	
	Priesley	104	104	106	106	111	
	Gallus (awned) 	99	101	101	102	107	
Means (t/ha)		5.80	5.91	6.20	6.17	6.62	
Means (bu/ac)		86	88	92	92	98	
Location-years		30	23	16	12	7	

¹ Values differing by less than 3 within a column may not represent true differences in yield. Yield Indices are Heritability Adjusted Relative Values (HARV), which favour results from trial locations with high repeatability. For more information, see: Yan, W. Use of HARV in Variety Trial Summaries.









² Area I and II combined: see map of areas on www.gocereals.ca/variety_trial.php.

³ sww = soft white winter, srw = soft red winter, hrw = hard red winter.

⁴ Cultivar yield rankings may vary from year to year. Decisions are therefore best made using data with the greatest number of years.

 PBR Status; indicates varieties protected under PBR 91. Visit pbrfacts.ca to learn more.

**Table 1a - Ontario Winter Wheat Performance Trial - Cumulative Yield Indices¹, Intensive trials
Summary for Area I & II Combined², OCCC, August 2016**

Class ³	Variety	5-Year Index Fungicides		4-Year Index Fungicides		3-Year Index Fungicides		2-Year Index Fungicides		2016 Index Fungicides	
		No	Yes	No	Yes	No	Yes	No	Yes	No	Yes
sww	Venture	86 ⁴	100	85	100	84	99	79	97	71	98
	25W31 (awned) 							96	104	92	100
srw	Emmit	95	105	95	105	92	103	90	104	89	103
	Branson	102	109	101	109	101	108	102	108	103	105
	Wave	97	104	97	102	95	100	95	101	98	102
	CM614	99	108	99	107	100	106	101	106	99	104
	WB425 (awned) 	99	105	99	105	98	104	99	104	99	102
	CM249 (awned)	101	108	100	108	99	106	99	107	103	108
	OAC Flight (awned) 	94	106	94	105	93	104	88	101	85	102
	25R34 (awned)	104	113	104	112	103	110	102	108	104	107
	25R40 (awned)	105	113	104	112	104	109	104	109	106	107
	HY 412-SRW 	97	106	97	106	96	104	97	103	95	100
	Emperor 	98	106	98	106	97	105	96	106	97	105
	Marker			100	109	102	109	101	108	103	108
	UGRC Ring (awned)			100	111	102	110	102	109	107	109
	UGRC C2-5					97	101	95	100	93	100
	25R46 (awned)							88	104	85	104
	Cruze (awned)							100	106	103	107
	DS572SRW 							99	110	100	111
	25R39									103	108
	Drew (awned)									94	104
	Curtis 									90	102
hrw	AC Morley	93	100	93	101	92	98	92	98	88	90
	AC Sampson	90	97	90	97	92	96	90	96	89	95
	Princeton	93	103	93	103	92	101	91	100	95	100
	Priesley	99	105	99	105	101	104	100	103	103	104
	Gallus (awned) 	94	100	96	101	97	100	97	102	102	104
Means (t/ha)		5.68	6.25	5.66	6.23	5.98	6.48	5.85	6.39	6.45	6.92
Means (bu/ac)		84	93	84	93	89	96	87	95	96	103
Location-years		22		19		13		9		5	

¹ Values differing by less than 3 within a column may not represent true differences in yield. Yield Indices are Heritability Adjusted Relative Values (HARV), which favour results from trial locations with high repeatability. For more information, see: Yan, W. Use of HARV in Variety Trial Summaries.


² Area I and II combined: see map of areas on www.gocereals.ca/variety_trial.php.

³ sww = soft white winter, srw = soft red winter, hww = hard white winter, hrw = hard red winter.









⁴ Cultivar yield rankings may vary from year to year. Decisions are therefore best made using data with the greatest number of years.

Fungicide treatments included a T1 application of foliar fungicide at Zadoks Growth Stage 30 to 31, followed by a T3 application of fusarium fungicide during flowering.

The OCCC is thankful to the Grain Farmers of Ontario, BASF, Bayer and Syngenta for their financial support towards the intensive trials.

 PBR Status; indicates varieties protected under PBR 91. Visit pbrfacts.ca to learn more.

**Table 2 - Ontario Winter Wheat Performance Trial; Cumulative Yield Index¹
Summary for Area I², OCCC, August 2016**


Class ³	Variety	5 year	4 year	3 year	2 year	2016
sww	Venture	91 ⁴	89	86	82	70
	25W31 (awned) 				101	97
srw	Emmit	98	98	95	92	88
	Branson	104	103	104	105	106
	Wave	101	98	95	95	94
	CM614	105	104	103	104	102
	WB425 (awned) 	103	101	102	104	104
	CM249 (awned)	107	106	104	104	105
	OAC Flight (awned) 	96	96	94	88	79
	25R34 (awned)	111	110	109	110	113
	25R40 (awned)	112	110	111	112	117
	HY 412-SRW 	101	100	98	97	98
	Emperor 	99	99	99	97	97
	Marker		103	104	103	107
	UGRC Ring (awned)			106	107	107
	UGRC C2-5				96	94
	25R46 (awned)				98	93
	Cruze (awned)					108
	DS572SRW 					100
	25R39					112
	Drew (awned)					
	Curtis 					
hrw	AC Morley	96	95	93	92	85
	AC Sampson	92	92	92	90	88
	Princeton	97	98	96	95	94
	Priesley	104	104	104	107	113
	Gallus (awned) 	96	99	99	101	105
Means (t/ha)		5.60	5.93	6.16	5.98	6.24
Means (bu/ac)		83	88	92	89	93
Location-Years		13	10	7	5	3

¹ Values differing by less than 3 within a column may not represent true differences in yield. Yield Indices are Heritability Adjusted Relative Values (HARV), which favour results from trial locations with high repeatability. For more information, see: Yan, W. Use of HARV in Variety Trial Summaries.









² Area I : see map of areas on www.gocereals.ca/variety_trial.php.

³ sww = soft white winter, srw = soft red winter, hrw = hard red winter.

⁴ Cultivar yield rankings may vary from year to year. Decisions are therefore best made using data with the greatest number of years.

 PBR Status; indicates varieties protected under PBR 91. Visit pbrfacts.ca to learn more.

**Table 2a - Ontario Winter Wheat Performance Trial - Cumulative Yield Indices¹, Intensive trials
Summary for Area I², OCCC, August 2016**

Class ³	Variety	5-Year Index Fungicides		4-Year Index Fungicides		3-Year Index Fungicides		2-Year Index Fungicides		2016 Index Fungicides	
		No	Yes	No	Yes	No	Yes	No	Yes	No	Yes
sww	Venture	89 ⁴	103	90	102	88	102	84	102	75	100
	25W31 (awned) 							100	108	94	100
srw	Emmit	95	104	95	104	92	102	88	103	88	103
	Branson	100	107	99	105	99	106	100	107	103	105
	Wave	98	104	96	102	93	99	92	98	94	96
	CM614	100	108	100	107	98	105	99	106	95	101
	WB425 (awned) 	96	104	96	104	97	106	98	107	96	100
	CM249 (awned)	103	109	102	107	100	106	99	105	101	105
	OAC Flight (awned) 	95	107	94	106	94	105	87	102	78	96
	25R34 (awned)	106	115	105	113	103	110	103	111	107	106
	25R40 (awned)	107	115	105	112	105	111	105	110	110	110
	HY 412-SRW 	96	104	96	103	94	100	91	99	93	99
	Emperor 	95	104	94	103	94	103	91	103	94	103
	Marker			99	106	100	107	98	109	107	108
	UGRC Ring (awned)			101	108	101	109	101	110	106	106
	UGRC C2-5					92	99	88	97	87	96
	25R46 (awned)					97	108	92	107	88	106
	Cruze (awned)							101	108	105	110
	DS572SRW 							96	111	93	108
25R39									106	109	
Drew (awned)									95	110	
Curtis 									93	104	
hrw	AC Morley	93	99	93	99	91	98	91	99	83	87
	AC Sampson	89	95	89	96	89	95	86	95	88	96
	Princeton	92	99	94	101	91	98	89	98	87	92
	Priesley	99	105	99	105	100	104	103	105	112	109
	Gallus (awned) 	92	98	94	100	95	101	95	102	102	106
Means (t/ha)	5.61	6.14	5.74	6.27	5.91	6.44	5.54	6.12	5.61	5.99	
Means (bu/ac)	83	91	85	93	88	96	82	91	83	89	
Location-Years		11		9		7		4		2	

¹ Values differing by less than 3 within a column may not represent true differences in yield. Yield Indices are Heritability Adjusted Relative Values (HARV), which favour results from trial locations with high repeatability. For more information, see: Yan, W. Use of HARV in Variety Trial Summaries.


² Area I: see map of areas on www.gocereals.ca/variety_trial.php.

³ sww = soft white winter, srw = soft red winter, hww = hard white winter, hrw = hard red winter.










⁴ Cultivar yield rankings may vary from year to year. Decisions are therefore best made using data with the greatest number of years.

Fungicide treatments included a T1 application of foliar fungicide at Zadoks Growth Stage 30 to 31, followed by a T3 application of fusarium fungicide during flowering.

The OCCC is thankful to the Grain Farmers of Ontario, BASF, Bayer and Syngenta for their financial support towards the intensive trials.

 PBR Status; indicates varieties protected under PBR 91. Visit pbrfacts.ca to learn more.

**Table 3 - Ontario Winter Wheat Performance Trial; Cumulative Yield Index¹
Summary for Area II², OCCC, August 2016**


Class ³	Variety	5 year	4 year	3 year	2 year	2016
sww	Ava	105 ⁴	105	106	106	105
	Venture	87	84	81	77	66
	25W31 (awned) 				96	91
srw	Emmit	101	100	96	94	91
	Branson	107	108	107	108	109
	Wave	102	103	101	101	102
	CM614	103	104	106	107	108
	WB425 (awned) 	105	106	104	103	106
	CM249 (awned)	104	104	103	104	109
	OAC Flight (awned) 	96	94	90	88	83
	25R34 (awned)	107	109	107	106	110
	25R40 (awned)	107	108	107	109	110
	HY 412-SRW 	102	103	103	104	100
	Emperor 	103	104	100	100	98
	Marker		106	107	106	108
	UGRC Ring (awned)		104	106	106	111
	UGRC C2-5			102	101	97
	25R46 (awned)			87	87	82
	Cruze (awned)				103	106
	DS572SRW 				103	104
	25R39				107	107
	Drew (awned)					91
	Curtis 					85
hrw	AC Morley	96	96	93	93	90
	AC Sampson	95	95	96	95	90
	Princeton	99	98	99	100	107
	Priesley	105	105	108	105	109
	Gallus (awned) 	102	103	103	103	109
	HY 301-HRW 	99	99	100	98	104
Means (t/ha)		5.95	5.89	6.23	6.30	6.88
Means (bu/ac)		88	88	93	94	102
Location-Years		17	13	9	7	4

¹ Values differing by less than 3 within a column may not represent true differences in yield. Yield Indices are Heritability Adjusted Relative Values (HARV), which favour results from trial locations with high repeatability. For more information, see: Yan, W. Use of HARV in Variety Trial Summaries.










² Area II : see map of areas on www.gocereals.ca/variety_trial.php.

³ sww = soft white winter, srw = soft red winter, hrw = hard red winter.

⁴ Cultivar yield rankings may vary from year to year. Decisions are therefore best made using data with the greatest number of years.

 PBR Status; indicates varieties protected under PBR 91. Visit pbrfacts.ca to learn more.

**Table 3a - Ontario Winter Wheat Performance Trial - Cumulative Yield Indices¹, Intensive Trials
Summary for Area II², OCCC, August 2016**

Class ³	Variety	5-Year Index Fungicides		4-Year Index Fungicides		3-Year Index Fungicides		2-Year Index Fungicides		2016 Index Fungicides	
		No	Yes	No	Yes	No	Yes	No	Yes	No	Yes
sww	Ava	101 ⁴	109	100	109	103	109	104	108	101	105
	Venture	83	98	81	98	80	96	75	93	69	97
	25W31 (awned) 							93	101	90	99
srw	Emmit	95	106	95	107	93	105	91	106	90	103
	Branson	103	110	103	111	103	110	104	109	104	105
	Wave	97	103	98	103	97	101	98	104	100	107
	CM614	98	107	98	108	101	106	102	107	102	107
	WB425 (awned) 	101	106	102	105	99	102	99	101	100	103
	CM249 (awned)	99	108	98	108	98	105	99	108	104	110
	OAC Flight (awned) 	94	105	94	105	92	102	90	100	90	106
	25R34 (awned)	102	112	103	112	102	109	100	106	103	107
	25R40 (awned)	103	111	103	111	103	108	104	109	103	104
	HY 412-SRW 	97	108	98	108	99	107	101	107	96	101
	Emperor 	100	108	101	108	99	107	99	108	99	106
	Marker			101	111	103	110	102	107	101	107
	UGRC Ring (awned)			99	113	102	111	102	109	108	111
	UGRC C2-5					101	102	101	103	97	103
	25R46 (awned)							85	102	83	103
	Cruze (awned)							99	104	102	105
	DS572SRW 							102	109	104	112
25R39									101	106	
Drew (awned)									93	100	
Curtis 									88	101	
hrw	AC Morley	93	101	92	102	92	99	92	98	91	92
	AC Sampson	92	98	91	97	94	96	93	97	89	95
	Princeton	93	105	92	105	93	104	92	101	99	106
	Priesley	99	105	99	105	101	104	97	102	97	102
	Gallus (awned) 	96	102	97	101	99	99	98	102	103	103
	HY 301-HRW 	94	96	94	96	95	95	93	97	98	99
Means (t/ha)	5.74	6.34	5.59	6.19	6.04	6.52	6.08	6.58	6.98	7.49	
Means (bu/ac)	85	94	83	92	90	97	90	98	104	111	
Location-Years	12		10		7		5		3		

¹ Values differing by less than 3 within a column may not represent true differences in yield. Yield Indices are Heritability Adjusted Relative Values (HARV), which favour results from trial locations with high repeatability. For more information, see: Yan, W. Use of HARV in Variety Trial Summaries.


² Area II: see map of areas on www.gocereals.ca/variety_trial.php.

³ sww = soft white winter, srw = soft red winter, hww = hard white winter, hrw = hard red winter.









⁴ Cultivar yield rankings may vary from year to year. Decisions are therefore best made using data with the greatest number of years.

Fungicide treatments included a T1 application of foliar fungicide at Zadoks Growth Stage 30 to 31, followed by a T3 application of fusarium fungicide during flowering.

The OCCC is thankful to the Grain Farmers of Ontario, BASF, Bayer and Syngenta for their financial support towards the intensive trials.

 PBR Status; indicates varieties protected under PBR 91. Visit pbrfacts.ca to learn more.

**Table 4 - Ontario Winter Wheat Performance Trial; Cumulative Yield Index¹
Summary for Area III², OCCC, August 2016**


Class ³	Variety	5 year	4 year	3 year	2 year	2016
sww	Ava	103 ⁴	103	103	104	104
srw	Emmit	103	103	102	103	110
	Branson	103	102	102	101	103
	CM614	106	108	108	109	114
	WB425 (awned) 	99	99	97	96	88
	CM249 (awned)	102	103	103	102	103
	HY 412-SRW 	102	104	102	103	102
	Emperor 	97	96	96	95	96
	OAC Flight (awned) 	98	95	95	95	92
	Marker		104	104	105	106
	UGRC Ring (awned)		107	107	108	113
	UGRC C2-5				105	105
	Cruze (awned)				96	91
	DS572SRW 				105	111
	Drew (awned)					97
Curtis 					106	
hrw	AC Morley	95	97	97	97	100
	AC Sampson	95	95	94	92	90
	Princeton	95	96	97	97	102
	Gallus (awned) 	99	98	96	95	97
	Priesley	103	102	102	99	96
	HY 301-HRW 	94	92	93	93	88
Means (t/ha)		5.43	4.99	4.67	4.39	4.22
Means (bu/ac)		81	74	69	65	63
Location-Years		11	8	7	5	2

¹ Values differing by less than 3 within a column may not represent true differences in yield. Yield Indices are Heritability Adjusted Relative Values (HARV), which favour results from trial locations with high repeatability. For more information, see: Yan, W. Use of HARV in Variety Trial Summaries.

² Area III : see map of areas on www.gocereals.ca/variety_trial.php.

³ sww = soft white winter, srw = soft red winter, hrw = hard red winter.

⁴ Cultivar yield rankings may vary from year to year. Decisions are therefore best made using data with the greatest number of years.

 PBR Status; indicates varieties protected under PBR 91. Visit pbrfacts.ca to learn more.

**Table 5a - Ontario Winter Wheat Performance Trials; Varietal Characteristics Based on Data from Area I
OCCC, August 2016**

Variety	Fusarium Data			Test Weight (kg/hL)	Protein (%)	1000 Kernel Weight (g)	Winter Survival (%)	Lodging (0-9) ²	Height (cm)	Heading Date ³ (JD)	Powdery Mildew (0-9) ²	Leaf Septoria (0-9) ²	Stripe Rust (0-9) ²	Eyespot (0-9) ²
	Combined Fusarium Rating ¹	Don Rating ¹	Years											
Venture	S	S	5	72.6	10.4	28.7	99	2.4	74	151	1.0	1.0	7.5	2.0
25W31	MS	MS	1	76.5	11.4	29.4	97	1.4	83	152	5.5	0.0	4.1	2.0
Emmit	MS	MS	7	76.5	10.4	35.2	97	1.3	86	152	5.0	0.0	4.3	2.8
Branson	S	MS	7	76.7	10.8	35.4	95	2.8	88	148	4.5	0.0	1.3	2.3
25R39	MS	MS	6	78.1	9.9	38.1	98	3.9	86	150	5.0	1.5	0.9	2.3
Wave	S	MS	7	78.6	11.2	34.1	97	3.1	98	150	5.0	0.0	4.3	3.0
CM614	MS	MS	6	78.9	10.4	38.7	97	5.6	94	149	2.5	0.0	3.8	6.0
WB425	MS	MR	5	78.2	10.4	31.9	98	1.1	86	148	0.5	0.0	2.7	2.8
CM249	S	MS	5	76.1	10.3	36.7	98	2.9	91	149	5.0	0.5	3.5	4.5
25R34	MR	MR	4	76.4	10.1	37.3	97	2.0	85	150	5.0	3.0	1.9	2.5
25R40	HS	S	4	78.2	10.1	38.9	92	1.0	76	150	1.5	1.0	1.5	0.8
HY 412-SRW	MS	S	4	78.9	11.1	37.8	93	1.5	88	153	3.0	0.0	2.9	1.5
Emperor	MS	MR	4	78.1	10.6	35.8	98	2.6	95	148	5.0	2.0	4.8	3.3
OAC Flight	MS	MS	5	76.2	11.0	32.8	91	1.5	84	151	4.0	0.5	4.7	1.8
Marker	MR	MR	3	76.4	9.9	32.0	98	1.3	86	151	3.0	0.5	3.2	1.8
UGRC Ring	MS	MS	3	76.7	10.1	37.1	99	2.4	93	150	5.0	1.5	3.1	3.3
UGRC C2-5	MR	MR	2	73.9	10.2	37.8	96	2.9	90	153	4.0	1.5	3.7	2.5
25R46	MR	MR	2	74.8	10.0	30.7	98	0.3	80	150	5.5	0.5	6.1	1.8
Cruze	MS	MS	1	78.6	10.7	34.8	96	1.3	84	149	5.0	2.0	3.1	1.3
DS572SRW	S	S	1	77.7	10.1	38.8	94	2.0	100	151	3.0	0.0	3.8	3.0
Drew	--	--	--	76.2	10.7	30.0	96	1.0	88	148	3.5	0.5	4.5	1.3
Curtis	--	--	--	76.8	10.2	31.2	97	3.0	91	150	2.0	0.5	5.6	1.5
AC Morley	MR	MR	7	78.6	12.0	35.4	99	4.4	114	151	3.0	3.0	3.9	5.3
AC Sampson	S	S	7	75.7	11.3	37.5	93	3.0	91	153	1.5	0.5	2.9	2.3
Princeton	MS	S	7	80.1	11.3	43.9	95	3.9	101	151	6.5	4.0	1.9	5.5
Priesley	S	MS	6	77.1	10.8	40.5	97	0.6	80	152	2.0	2.0	0.8	0.3
Gallus	MS	MS	5	81.4	11.6	42.8	95	1.4	89	150	4.0	1.5	1.1	1.5
Means				77.2	10.6	35.7	96	2.2	89	150	3.7	1.0	3.4	2.5
Locations				2	2	3	1	2	2	2	1	1	3	1

For traits not reported on this table see other area trait tables or previous year's performance trials. Not all traits are expressed in any given year or area. Head to Head comparisons on GoCereals.ca website give multi-year data for trait characteristics.

1 Combined Fusarium Ratings are based on BOTH Fusarium head blight ratings and deoxynivalenol (DON) levels from inoculated provincial trials.

DON Ratings are based only on DON levels from inoculated provincial trials.










MR=moderately resistant (best); MS=moderately susceptible; S=susceptible; HS=highly susceptible (worst).

2 For ratings 0-9, a high score is undesirable.

3 Heading may vary from year to year and should only be used to indicate relative differences.

 = PBR Status; indicates varieties protected under PBR 91. Visit pbrfacts.ca to learn more.

**Table 5b - Ontario Winter Wheat Performance Trials; Varietal Characteristics Based on Data from Area II
OCCC, August 2016**

Variety	Fusarium Data			Test Weight (kg/hL)	Protein (%)	1000 Kernel Weight (g)	Winter Survival (%)	Lodging (0-9) ²	Height (cm)	Heading Date ³ (JD)	Powdery Mildew (0-9) ²	Leaf Septoria (0-9) ²	Stripe Rust (0-9) ²
	Combined Fusarium Rating ¹	Don Rating ¹	Years										
Ava	MR	MR	7	78.3	8.7	36.3	90	0.3	101	157	3.0	5.3	3.6
Venture	S	S	5	74.5	9.7	28.4	89	0.0	78	157	0.0	7.0	6.6
25W31 	MS	MS	1	78.1	10.7	30.3	90	0.0	84	157	1.8	5.0	4.8
Emmit	MS	MS	7	77.0	9.5	35.9	91	0.5	91	157	3.5	6.0	4.6
Branson	S	MS	7	77.7	9.5	37.9	91	0.8	88	154	2.5	4.5	1.1
25R39	MS	MS	6	79.1	9.3	40.8	83	1.0	87	156	3.4	4.8	0.9
Wave	S	MS	7	78.9	9.8	34.9	94	1.3	95	155	2.5	5.8	4.5
CM614	MS	MS	6	79.4	9.2	37.7	97	2.0	93	156	2.3	5.0	2.8
WB425 	MS	MR	5	79.2	9.6	32.3	87	0.0	89	155	0.6	5.0	1.9
CM249	S	MS	5	77.2	9.3	39.2	94	0.8	95	155	2.6	6.0	3.2
25R34	MR	MR	4	77.7	9.3	40.0	91	0.3	87	155	3.3	4.5	1.5
25R40	HS	S	4	78.2	9.3	37.7	87	0.0	77	156	1.0	4.8	0.9
HY 412-SRW 	MS	S	4	80.3	10.0	37.5	90	0.0	91	159	1.7	4.5	2.7
Emperor 	MS	MR	4	77.4	9.6	32.8	97	1.8	100	155	3.4	5.5	6.1
OAC Flight 	MS	MS	5	75.4	9.7	34.6	90	0.0	87	156	2.4	6.5	6.4
Marker	MR	MR	3	76.3	9.3	32.1	90	0.3	92	156	2.4	4.0	1.8
UGRC Ring	MS	MS	3	76.8	9.0	37.5	95	0.5	96	156	3.2	6.3	3.5
UGRC C2-5	MR	MR	2	76.4	9.1	40.2	85	1.8	97	159	2.2	5.5	4.0
25R46	MR	MR	2	75.4	9.6	31.6	90	1.0	83	156	2.2	6.8	5.9
Cruze	MS	MS	1	79.3	9.8	35.1	90	2.0	85	155	3.0	6.3	2.7
DS572SRW 	S	S	1	78.9	9.2	42.4	97	0.3	102	156	0.0	5.3	3.4
Drew	--	--	--	77.8	9.8	30.1	85	0.0	86	155	1.7	6.3	4.5
Curtis 	--	--	--	77.2	9.6	30.0	88	0.3	91	156	0.3	6.5	5.7
AC Morley	MR	MR	7	79.8	10.5	36.9	94	2.8	115	157	2.0	6.0	2.4
AC Sampson	S	S	7	76.8	10.6	39.3	86	1.0	93	160	2.9	5.3	2.4
Princeton	MS	S	7	81.0	9.9	45.5	96	0.8	104	157	4.0	4.3	1.5
Priesley	S	MS	6	78.6	10.0	42.0	90	0.0	81	156	0.0	4.3	0.6
Gallus 	MS	MS	5	81.6	10.4	43.6	91	0.0	95	156	0.6	3.5	0.4
HY 301-HRW 	MS	MS	4	82.2	10.6	36.6	94	0.8	92	156	1.8	5.0	0.9
Means				78.2	9.7	36.5	91	0.7	91	156	2.1	5.3	3.1
Locations				4	2	4	3	1	4	4	2	1	4

For traits not reported on this table see other area trait tables or previous year's performance trials. Not all traits are expressed in any given year or area. Head to Head comparisons on GoCereals.ca website give multi-year data for trait characteristics.


1 Combined Fusarium Ratings are based on BOTH Fusarium head blight ratings and deoxynivalenol (DON) levels from inoculated provincial trials.

DON Ratings are based only on DON levels from inoculated provincial trials.









MR=moderately resistant (best); MS=moderately susceptible; S=susceptible; HS=highly susceptible (worst).

2 For ratings 0-9, a high score is undesirable.

3 Heading may vary from year to year and should only be used to indicate relative differences.

 = PBR Status; indicates varieties protected under PBR 91. Visit pbrfacts.ca to learn more.

**Table 5c - Ontario Winter Wheat Performance Trials; Varietal Characteristics Based on Data from Area III
OCCC, August 2016**

Variety	Fusarium Data			Test Weight (kg/hL)	Protein (%)	1000 Kernel Weight (g)	Winter Survival (%)	Height (cm)	Heading Date ³ (JD)	Leaf Septoria (0-9) ²
	Combined Fusarium Rating ¹	Don Rating ¹	Years							
Ava	MR	MR	7	75.2	11.0	31.8	80	80	158	2.7
Emmit	MS	MS	7	75.5	11.9	33.6	83	72	158	4.7
Branson	S	MS	7	75.2	12.5	33.4	77	66	155	2.7
CM614	MS	MS	6	75.6	12.2	33.6	81	74	159	4.7
WB425 	MS	MR	5	76.4	13.2	30.4	77	68	156	2.3
CM249	S	MS	5	74.0	12.4	34.9	78	72	156	3.3
HY 412-SRW 	MS	S	4	76.3	13.0	33.9	81	77	159	3.7
Emperor 	MS	MR	4	74.9	12.1	32.7	76	80	157	3.7
OAC Flight 	MS	MS	5	74.6	14.4	35.5	73	64	157	2.7
Marker	MR	MR	3	75.3	11.8	29.2	83	73	156	3.3
UGRC Ring	MS	MS	3	75.8	11.8	35.7	81	74	157	3.0
UGRC C2-5	MR	MR	2	75.3	12.3	38.0	75	75	162	3.3
Cruze	MS	MS	1	76.9	13.1	31.2	76	64	157	3.3
DS572SRW 	S	S	1	76.2	12.4	40.6	78	81	158	3.7
Drew	--	--	--	74.9	11.7	29.0	79	65	157	2.0
Curtis 	--	--	--	76.0	12.2	31.3	77	77	158	3.0
AC Morley	MR	MR	7	76.9	13.5	34.4	77	96	158	4.3
AC Sampson	S	S	7	73.3	14.1	35.0	79	75	163	3.0
Princeton	MS	S	7	76.5	13.2	38.7	79	77	158	2.7
Priesley	S	MS	6	71.8	13.5	35.0	79	61	158	2.7
Gallus 	MS	MS	5	76.9	14.5	38.7	80	69	159	5.3
HY 301-HRW 	MS	MS	4	77.0	14.6	31.5	79	68	158	4.0
Means				75.5	12.8	34.0	78	73	158	3.4
Locations				2	1	2	1	2	1	1

For traits not reported on this table see other area trait tables or previous year's performance trials. Not all traits are expressed in any given year or area. Head to Head comparisons on GoCereals.ca website give multi-year data for trait characteristics.

1 Combined Fusarium Ratings are based on BOTH Fusarium head blight ratings and deoxynivalenol (DON) levels from inoculated provincial trials.

DON Ratings are based only on DON levels from inoculated provincial trials.





MR=moderately resistant (best); MS=moderately susceptible; S=susceptible; HS=highly susceptible (worst).

2 For ratings 0-9, a high score is undesirable.

3 Heading may vary from year to year and should only be used to indicate relative differences.

 = PBR Status; indicates varieties protected under PBR 91. Visit pbrfacts.ca to learn more.

**Table 6 - Ontario Winter Wheat Performance Trial; Cumulative Yield Index¹
Summary for AreaV², OCCC, August 2016**


Class ³	Variety	3 year	2 year	2016
sww	Ava	104 ⁴	105	98
srw	CM614	97	98	98
	HY 412-SRW 	107	109	114
	UGRC C2-5		98	102
	DS572SRW 		105	98
	Curtis 			102
hrw	Princeton	95	94	93
	Priesley	105	102	104
	Gallus (awned) 	90	88	86
Means (t/ha)		7.99	7.99	7.52
Means (bu/ac)		119	119	112
Location-Years		4	3	1

¹ Values differing by less than 3 within a column may not represent true differences in yield. Yield Indices are Heritability Adjusted Relative Values (HARV), which favour results from trial locations with high repeatability. For more information, see: Yan, W. Use of HARV in Variety Trial Summaries.





² Area V : see map of areas on www.gocereals.ca/variety_trial.php.

³ sww = soft white winter, srw = soft red winter, hrw = hard red winter.

⁴ Cultivar yield rankings may vary from year to year. Decisions are therefore best made using data with the greatest number of years.

 PBR Status; indicates varieties protected under PBR 91. Visit pbrfacts.ca to learn more.

**Table 6a - Ontario Winter Wheat Performance Trial - Cumulative Yield Indices¹,
Intensive Trials Summary for Area V², OCCC, August 2016**

Class ³	Variety	3-Year Index Fungicides		2-Year Index Fungicides		2016 Index Fungicides	
		No	Yes	No	Yes	No	Yes
sww	Ava	104 ⁴	101	99	95	99	92
srw	CM614	97	99	95	97	99	99
	HY 412-SRW 	106	107	108	109	114	114
	UGRC C2-5					103	103
	DS572SRW 					99	96
	Curtis 					103	101
hrw	Princeton	98	99	95	98	94	94
	Priesley	104	103	107	106	105	99
	Gallus (awned) 	94	93	92	92	87	86
Means (t/ha)		7.35	7.40	7.77	7.84	7.52	7.37
Means (bu/ac)		109	110	116	117	112	110
Location-Years		3		2		1	

¹ Values differing by less than 3 within a column may not represent true differences in yield. Yield Indices are Heritability Adjusted Relative Values (HARV), which favour results from trial locations with high repeatability. For more information, see: Yan, W. Use of HARV in Variety Trial Summaries.


² Area V: see map of areas on www.gocereals.ca/variety_trial.php.

³ sww = soft white winter, srw = soft red winter, hww = hard white winter, hrw = hard red winter.





⁴ Cultivar yield rankings may vary from year to year. Decisions are therefore best made using data with the greatest number of years.

Fungicide treatments included a T1 application of foliar fungicide at Zadoks Growth Stage 30 to 31, followed by a T3 application of fusarium fungicide during flowering.

The OCCC is thankful to the Grain Farmers of Ontario, BASF, Bayer and Syngenta for their financial support towards the intensive trials.

 PBR Status; indicates varieties protected under PBR 91. Visit pbrfacts.ca to learn more.

**Table 6b - Ontario Winter Wheat Performance Trials; Varietal Characteristics Based on Data from Area V
OCCC, August 2016**

Variety	Fusarium Data			Test Weight (kg/hL)	Protein (%)	1000 Kernel Weight (g)	Lodging (0-9) ²	Height (cm)	Heading Date ³ (JD)
	Combined Fusarium Rating ¹	Don Rating ¹	Years						
Ava	MR	MR	7	82.3	11.7	35.3	0.5	109	171
CM614	MS	MS	6	82.1	12.0	38.7	1.8	103	170
HY 412-SRW 	MS	S	4	82.2	11.9	35.9	0.0	97	171
UGRC C2-5	MR	MR	2	79.3	11.5	39.5	0.3	106	170
DS572SRW 	S	S	1	82.2	12.0	44.7	2.8	108	168
Curtis 	--	--	--	82.3	12.2	39.5	2.5	102	168
Princeton	MS	S	7	82.8	12.9	45.2	0.0	106	170
Priesley	S	MS	6	82.2	12.4	42.5	0.0	92	173
Gallus 	MS	MS	5	83.5	13.9	44.6	0.0	97	168
Means				82.1	12.3	40.7	0.9	102	170
Locations				1	1	1	1	1	1

For traits not reported on this table see other area trait tables or previous year's performance trials. Not all traits are expressed in any given year or area. Head to Head comparisons on GoCereals.ca website give multi-year data for trait characteristics.

1 Combined Fusarium Ratings are based on BOTH Fusarium head blight ratings and deoxynivalenol (DON) levels from inoculated provincial trials.
DON Ratings are based only on DON levels from inoculated provincial trials.










MR=moderately resistant (best); MS=moderately susceptible; S=susceptible; HS=highly susceptible (worst).

2 For ratings 0-9, a high score is undesirable.

3 Heading may vary from year to year and should only be used to indicate relative differences.

 = PBR Status; indicates varieties protected under PBR 91. Visit pbrfacts.ca to learn more.

Table 7a - Ontario Winter Wheat Distributors

Class ³	Variety	Distributor
sww	Ava	Dow Seeds
	Venture	SeCan Association
	25W31 (awned) 	DuPont Pioneer
srw	Emmit	Dow Seeds
	Branson	Dow Seeds
	Wave	SeCan Association
	CM614	C & M Seeds
	WB425 (awned) 	C & M Seeds
	CM249 (awned)	C & M Seeds
	OAC Flight (awned) 	Elite Seeds
	25R34 (awned)	DuPont Pioneer
	25R40 (awned)	DuPont Pioneer
	HY 412-SRW 	Dow Seeds
	Emperor 	SeCan Association
	Marker	Bramhill Seeds
	UGRC Ring (awned)	Elite Seeds
	UGRC C2-5	Ridgetown Campus, U. of Guelph
	25R46 (awned)	DuPont Pioneer
	Cruze (awned)	C & M Seeds
DS572SRW 	Dow Seeds	
25R39	DuPont Pioneer	
Drew (awned)	C & M Seeds	
Curtis 	SeCan Association	
hrw	AC Morley	Advantage Seed Growers
	AC Sampson	Bramhill Seeds
	Princeton	C & M Seeds
	Priesley	C & M Seeds
	Gallus (awned) 	C & M Seeds
HY 301-HRW 	Dow Seeds	

¹ sww = soft white winter, srw = soft red winter, hrw = hard red winter.


 = PBR Status; indicates varieties protected under PBR 91. Visit pbrfacts.ca to learn more.

Table 7b - Ontario Winter Wheat Distributor/Sponsor Contact Information

Distributor	Address
Advantage Seed Growers	P.O. Box 351, Lucknow, ON, N0G 2H0 Tel: 519-440-6406 Fax: 519-812-1010 Internet: www.advantageseeds.com
Bramhill Seeds	5220 Hwy 23, RR#2, Palmerston, ON, N0G 2P0 Tel: 519-343-3630 Fax: 519-343-2037 Internet: www.bramhillseeds.com
C&M Seeds	6180 5th Line, Palmerston, ON, N0G 2P0 Tel: 1-888-733-9432 Fax: 519-343-3792 Internet: www.redwheat.com
Dow Seeds	P.O. Box 1090, 5 Hyland Drive Blenheim, ON, N0P 1A0 Tel: 519-676-8146 Fax: 519-676-6800 Internet: www.dowseeds.com
Elite Seeds	2405, rue de la Province, Longueuil, QC, J4G 1G3 Tel: 450-670-2231 Fax: 450-670-3900 Internet: www.eliteseeds.ca
DuPont Pioneer	Box 730, 7398 Queen's Line, Chatham, ON, N7M 5L1 Tel: 1-800-265-9435 Internet: www.pioneer.com
Ridgetown Campus, University of Guelph	120 Main St E, Ridgetown, ON, N0P 2C0 Tel: 1-877-674-1610 Internet: www.ridgetownc.uoguelph.ca
SeCan Association	400-300 Terry Fox Drive, Kanata, ON, K2K 0E3 Tel: 1-866-797-7874 Fax: 613-592-9497 Internet: www.secan.com